

US RENAL/BLADDER

Well hydrated.

Infants : give usual feeding 15 minutes before exam
Children : 8-16 oz. water upon arrival (no carbonated beverages)
Adolescents/young adults: 32 oz. water upon arrival (no carbonated beverages)

US PELVIS

Well hydrated.

Infants : give usual feeding 15 minutes before exam
Children : 8-16 oz. water upon arrival (no carbonated beverages)
Adolescents/young adults: 32 oz. water upon arrival (no carbonated beverages)

US ABDOMEN

Newborn to 4 months:	NPO 3 hours prior to exam
5 ó 8 months:	NPO 4 hours prior to exam
9 ó 12 months:	NPO 5 hours prior to exam
1 - 2 years :	NPO 6 hours prior to exam
.>2 years :	NPO 8 hours prior to exam

US ABDOMEN/PELVIS

Newborn to 4 months:	NPO 3 hours prior to exam
5 ó 8 months:	NPO 4 hours prior to exam
9 ó 12 months:	NPO 5 hours prior to exam
1 - 2 years :	NPO 6 hours prior to exam
>2 years :	NPO 8 hours prior to exam

NPO for 2 hours prior to exam.

For potty trained patients, advise to arrive with full bladder if possible, and not to empty bladder until pelvic portion of the examination is completed. If the child cannot cooperate, the abdominal portion of the examination will be completed and following that, the patient will be advised by the sonographer as to how much to drink to fill the bladder as quickly as possible so that pelvic portion of the examination can be performed.

US PYLORUS

NPO 3 hours prior to exam.

US INTUSSUSCEPTION

No prep.

US FOR APPENDICITIS

Advise patient not to void if potty trained.

Infants – 10 years : 2-16 oz water as tolerated

10 years – young adult: 32 oz water as tolerated.

US INFANT HIP FOR DEVELOPMENTAL DYSPLASIA (DDH)

NPO 3 hours prior to exam.

Bring along bottle of breast milk or formula to be fed to the baby during the sonogram so that infant is calm during the exam.

US HIP FOR EFFUSION

No prep.

US SPINE

Feed infant 30 minutes prior to examination.

US HEED i

BE

If diagnosis is for Hirschsprung's disease:

No prep.

For 2 days prior to BE:

- no rectal exams
- no rectal stimulation
- no rectal thermometers
- no suppositories
- no enemas

If diagnosis is not Hirschsprung's disease, stricture, pre-op for reanastomosis, intussusception, or constipation:

Speak with pediatric radiologist before scheduling. Prep will be ordered as per pediatric radiologist.

CONTRAST ENEMA FOR INTUSSUSCEPTION

No prep.

VCUG

NPO 1 hour prior to exam.

Fistulogram 3.87Tm0.502 g0.502 G[p]11(r)-6-4(o)11(r)-6 11.04 Tf1 0 0 1